



Playing the hand we're dealt

That <u>one population</u> on exceptionally <u>poor habitat</u> did not show a significant density-dependent response for recruitment may suggest that <u>some habitats are too poor for such a process to be operative or detectable.</u>


KEYSER, P.D., C.G. GUYNN, JR., AND H.S. Hill, JR. 2005 Density-dependent recruitment patterns in white-tailed deer. Wildlife Society Bulletin 33(1):222-232

... density independent factors such as rainfall and hard mast exert tremendous influence on these herds through their effect on nutrition. In the absence of either rain or mast, baseline levels of nutrition are simply too poor to allow for any alteration in either herd dynamics or condition. The habitat at <u>BR</u> is of such <u>low quality</u> that such a pattern would be expected (Shea and Osborne 1995)

KEYSER, P.D., C.G. GUYNN, JR., AND H.S. Hill, JR. 2005. Population density-physical condition relationships in white-tailed deer. Journal of Wildlife Management 69:356-365



1999 – 4,500 ac.



2000 Hunter Survey

51% thought the best management strategy for buck harvest was mandatory antler restrictions in specific areas, but not in all hunting areas.

44% thought that antier restrictions would hinder efforts to introduce young hunters to deer hunting. 53% thought they would help


56% would support antier restrictions for all hunting areas on Fort Bragg. 39% would oppose antier restrictions for all hunting areas on Fort Bragg.

Ginal Dack


Quality Deer Management is a popular term used today. The word "quality" means different things to different people, Which of the following characteristics do you consider when defining "quality deer"?

- a. Sex of the deer
- b. Age of the deer
- c. Weight of the deer
- d. Number of Points

- e. Antler spread
- f. Antler mass
- g. None of the above




Which one of the above characteristics BEST defines the term "Quality Deer"?






2000 - 13,000 ac.






2001 - 22,000 ac.

















Doe Harvest




(Grold) back.)



Antlered Buck Age Structure QDM (1999-2008)




Antlered Buck Age Structure Pre-QDM (1989-1997)






Mean Weights 1999-2008 (kg)







Mean Inside Spread (in)






Mean Number of Points 1999-2008








Mean Beam Diameters 1999-2008 (in)






Antlered Bucks by Mgmt. Area 2001-2008






Antlered Bucks Harvested/Mile2





QDM vs. Post Antlered Age Class Structure (1999-2008)



























Future Direction

- •Follow-up Hunter Survey 2009-2010
- Agricultural Outlease plan developed for area
- Possible expansion or additional management areas with antler restrictions
- Broad-based support, hunter buy-in, promising results in spite of habitat limitations



